

## About Corteva Agriscience™

- A global leader in seed and crop protection created from the former agricultural businesses of Dow AgroSciences, DuPont and Pioneer
- A strong portfolio comprising grassland & maize crop protection, silage inoculants and maize seed
- Corteva's significant investment in innovative science to find and develop new solutions is helping livestock farmers achieve their grassland and forage crop potential

**Use plant protection products safely. Always read the label and product information before use. Pay attention to the risk indications and follow the safety precautions on the label.**

**Triple rinse containers and invert to dry at time of use.**

Authorisation Holder: Corteva Agriscience, CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE. Tel: 01462 457272.

Marketing Company: Whelehan Crop Protection, Suite 11/12 Bunkilla Plaza, Bracetown Business Park, Clonee, Co. Meath, Ireland. Tel: 01 4688900 Email: [cropprotection@tpwhelehan.ie](mailto:cropprotection@tpwhelehan.ie)

®, ™ Trademarks of Corteva Agriscience and its affiliated companies.  
© 2021 Corteva. Supercedes all previous versions of this leaflet.

Forefront® T contains aminopyralid and triclopyr.



For use with  
a boom  
sprayer



**Forefront T** delivers the highest levels of control of docks, thistles, nettles, ragwort, buttercups and dandelions.

It can only be used on grazing ground or after last cut of silage. Manure restrictions apply.

**Seek advice before use.  
See product label for full details.**



To download the Corteva Grassland App, visit your device App Store and search for "Corteva Grassland".

You need to register the app on each individual device. The desktop version for Ireland is available at:

<https://grassland.farming.co.uk/#/mainMenu>



**For grassland advice call Whelehan Crop Protection on: 01 4688900 or visit: [www.corteva.ie/grassland](http://www.corteva.ie/grassland) or email: [cropprotection@tpwhelehan.ie](mailto:cropprotection@tpwhelehan.ie)**

### \*Ragwort label guidance

Where ragwort is present users should consult the Code of Practice on How to Prevent the Spread of Ragwort. Ragwort plants sprayed with this herbicide are more palatable and contain higher levels of toxins. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visible sign of the dead weed. Do not include treated ragwort in hay or silage crops.

# Forefront® T

## HERBICIDE

# Better weed control, better grazing.

grow great grass

grow great grass



# Forefront® T



## HERBICIDE

The widest spectrum and most effective herbicide available to grassland farmers. It is very safe to grass.

### Control grassland weeds because:

- They compete with grass for space, light, nutrients and water
- They are unpalatable to stock
- Thistles can lead to a greater incidence of Orf
- Ragwort poses a serious threat to livestock health and reduces grazing.

### Choose Forefront T because it:

- Is the best available weed control solution in cattle and/or sheep grazed grassland
- Can also be used on silage ground after the last cut of the year
- It moves to the roots ensuring high levels of long-term weed control
- Gives excellent grass safety.

Thistle and dock population can be calculated by counting the number of weeds in a 5 x 7m block. One weed will represent 1% weed infestation.



SAC – trials data from the Scottish Agricultural College shows 10% weed infestation causes 10% YIELD LOSS

### Key points:

<b>Product</b>	<b>Forefront T</b>
<b>Active ingredients</b>	30 g/L aminopyralid + 240 g/L triclopyr
<b>Weeds controlled</b>	
<b>Application rate</b>	2.0 L/ha
<b>Water volume</b>	200lt – 300lt/ha for high weed numbers or dense grass swards
<b>Maximum total dose</b>	2.0 L/ha per year
<b>Weed size</b>	
Broadleaved dock and curled dock	Rosette stage, 150 to 250mm across or high
Creeping thistle and spear thistle	Rosette stage, 150 to 250mm across or high
Common nettle	Actively growing, up to 300mm high
Creeping buttercup and dandelion	Actively growing, before flowering
Ragwort	Rosette stage, up to 200mm across or high
<b>Weed health</b>	Weeds must be actively growing; free from disease or insect damage; not suffering from drought, waterlogging or nutrient deficiency
<b>Stock exclusion</b>	7 days – only use on grazing ground grazed by cattle or sheep*
<b>Cutting interval</b>	Do not use Forefront T on fields to be cut for silage, hay or haylage unless it is after the last cut
<b>Manure</b>	If manure is generated, keep on farm and apply to grazing grassland
<b>Rainfastness</b>	1 hour when applied to a dry leaf
<b>Clover</b>	Will be damaged; re-introduce after 4 months

### Weeds controlled by Forefront T

Where we have knowledge of how Forefront T might affect other grassland weeds we have detailed it in the following tables. These are not recommendations; just an indication of what control might be achieved. ▽ indicates information based on anecdotal or limited data, and as such the user bears the risk in respect of failures concerning efficacy and phytotoxicity.

#### Annual weeds

Bindweed (black)	Fool's parsley	Orache
Bindweed (field)	Forget-me-not	Pale persicaria
Bristly ox-tongue	Fumitory	Poppy
Charlock	Groundsel	Redshank
Chickweed	Hemp-nettle	Scarlet pimpernel
Cleavers	Himalayan balsam	Shepherd's-purse
Corn chamomile	Knotgrass (4TL)	Speedwells
Corn marigold	Mayweeds	Spurrey
Cranesbill	Nettle (small)	Wild radish
Dead-nettles	Nightshade (black)	Yellow rattle
Fat-hen (2TL)		

#### Perennial weeds

Bramble	Ground elder	Old man's beard
Broom	Ground ivy	Plantain (greater)
Burdock	Hawthorn	Plantain (ribwort)
Buttercups	Hemlock	Ragwort
Cinquefoil	Hogweed (giant)	Rosebay willowherb
Clover, trefoil	Horsetail ( <i>Equisetum</i> )	Rushes
Coltsfoot	Japanese knotweed	Self-heal
Cow parsley	Knapweed (common)	Silverweed
Daisy (common)	Lesser celandine	Sorrel (common)
Daisy (ox-eye)	Mallow	Thistles
Dandelion	Mugwort	Vetch, tare
Docks	Nettle (common)	Yarrow
Gorse	Medick	Yellow/Flag Iris

<b>Weed control key</b>		No control
Good control		No information
Moderate control	▽	Anecdotal or limited information
Some control	TL	= true leaves