

PCS05808



IRELAND TECHNICAL INFORMATION SHEET

A broad spectrum herbicide for
**Established Grassland, New Sown Leys,
 Grass for Seed, and Maize**

Key Facts

Product Registration Number:	PCS 05808
Active Ingredient:	100 g / litre fluroxypyr + 80 g / litre clopyralid + 2.5 g / litre florasulam
Pack Size:	3 litre
Formulation:	Emulsifiable Concentrate
Maximum Individual Dose:	2.0 L/h to established grassland 1.0 L/ha to new sown leys, grass for seed and maize
Maximum Total Dose:	2.0 L/h to established grassland 1.0 L/ha to new sown leys, grass for seed and maize
Timing:	1 st February to 30 th September for established grassland and grass for seed 1 st February to 31 st August for new sown leys 10 th April to 31 st May (BBCH 13 – 16) for maize
Water Volumes:	200 - 400 L/ha
Spray Quality:	MEDIUM as defined by BCPC

Key Benefits

- 3 actives delivering effective translocation and wide spectrum of weed control
- Controls many weeds such as chickweed, fat-hen, mayweeds, buttercups, docks and thistles
- Convenient size pack which at 2 L/ha treats 1.5 ha (3.7 acres)
- Good grass safety
- Great versatility with use in established grassland, new sown leys, grass grown for seed and maize
- Rainfast in 2 hours

Best Use Advice - Grass

- For established grassland use at 2 L/ha to get excellent control of docks, thistles and buttercups.
- If weeds grow beyond the ideal stage, consider cutting the sward and spraying the regrowth
- Allow 2-3 weeks after cutting before applications of PastorTrio are made to allow sufficient regrowth
- For new sown leys apply at 1 L/ha to control a very wide range of broad-leaved weeds. Will kill clover
- Broad-leaved weeds not present at application will not be controlled
- Allow an interval of 4 weeks after application before cutting grass to optimise the effect of the herbicide
- Do not roll or harrow grass for 10 days before or 7 days after application
- Grass seed can be stitched in 1 month after application
- Clover can be stitched in 3 months after application
- Grazing can resume just 7 days after application
 - This may need to be longer if foliage of poisonous weeds such as ragwort is present
- To avoid subsequent injury to crops other than grassland and cereals, all spraying equipment must be thoroughly cleaned using All Clear Extra spray cleaner

Best Use Advice - Maize

- Apply when forage maize is between 3-6 leaf stage, and before the crop is over 20 cm tall. Do not apply once the buttress roots have started to develop on the first node.
- Crops that can be sown in the year of treatment with Leystar to forage maize include:
 - Cereals, oilseed rape, grass and vegetable brassicas as transplants
- To protect surface water, do not apply to maize before 10th April
- To avoid subsequent injury to crops other than grassland and cereals, all spraying equipment must be thoroughly cleaned using All Clear Extra spray cleaner